

The Ptolemaic Gamma-Ray Burst Universe

J. I. Katz

Department of Physics and McDonnell Center for the Space Sciences

Washington University, St. Louis

arXiv:astro-ph/9204003v2 30 Apr 1992

2 Copies, 10 pp., 0 figures, 0 tables

Mailing address: Department of Physics, Washington University, St. Louis, Mo. 63130

Telephone: 314-935-6202

e-mail: katz@wuphys.wustl.edu

Abstract

The BATSE experiment on GRO has demonstrated the isotropic arrival directions and flat $\log N$ *vs.* $\log S$ of cosmic gamma-ray bursts. These data are best explained if the burst sources are distributed throughout an extended spherical Galactic halo, as previously suggested by Jennings. The halo's radius is at least 40 Kpc, and probably is more than 100 Kpc. I consider possible origins of this halo, including primordial formation and neutron stars recoiling from their birthplaces in the Galactic disc. A simple geometrical model leads to a predicted relation between the dipole and quadrupole anisotropy. I suggest that neutron stars born with low recoil become millisecond pulsars, while those born with high recoil become the sources of gamma-ray bursts; these populations are nearly disjoint. Quiescent counterparts of gamma-ray bursts are predicted to be undetectably faint.

The first results from the BATSE on GRO (BATSE Science Team 1991) have revived the question of the distribution of gamma-ray burst sources in space. Their chief results, isotropy of gamma-ray burst directions and a $\log N$ *vs.* $\log S$ slope significantly flatter than -1.5, confirm earlier reports (see, for example, Meegan, Fishman and Wilson 1985 and the review by Cline 1984). Questions of relative calibration of different instruments and the paucity of good directional data permitted skepticism in the past. Such skepticism is no longer tenable, and the theoretical questions raised earlier must be faced.

An isotropic distribution of sources implies that, out to the maximum distance of observation permitted by instrumental sensitivity, all directions contain equivalent source populations. The source population for an observed flux or fluence S is expressed as the integral

$$N(\hat{\Omega}, S) = \int n(\hat{\Omega}, r, 4\pi r^2 S) 4\pi r^2 dr, \quad (1)$$

where $n(\hat{\Omega}, r, 4\pi r^2 S)$ is the density of sources at a distance r from the observer in the direction $\hat{\Omega}$ radiating a total power or energy $4\pi r^2 S$; a temporal average is implicit, and it is assumed that the sources radiate isotropically. While it is possible to obtain N independent of $\hat{\Omega}$ for a variety of artificial spatial distributions n , the only plausible n which does not require the observer to be in a preferred position (for example, at the center of spherical shells) is a homogeneous distribution out to the greatest distances at which the most luminous burst may be observed. Such a distribution implies $N \propto S^{-3/2}$, in contradiction to observation. This dilemma was fully appreciated and discussed by several of the contributors to the 1983 Symposium on High Energy Transients in Astrophysics (Woosley 1984).

The only geometric resolution of this problem is the Ptolemaic gamma-ray burst universe, as discussed by Cline (1984). The flat dependence of N on S (equivalent to a peak in a V/V_{max} distribution at $V \ll V_{max}$, and to a deficiency of faint bursts) is explained if the sources are distributed throughout a volume of finite extent, with most bursts within the volume luminous enough to be observed. This volume is spherical, with

the Earth at its center. The sources must be at cosmological distances, or be distributed in an extended halo around our Galaxy (such as the Massive Extended Halo suggested by Jennings 1984), or be evidence of a previously unsuspected Earth-centered spherical distribution.

At cosmological distances a typical burst of 10^{-7} erg/cm² fluence must have radiated $\sim 10^{50}$ erg in gamma-rays. This is energetically consistent with the spiraling-in or coalescence of binary neutron stars, if they convert gravitational energy to gamma-rays, perhaps through a magnetohydrodynamic dynamo, or with a relativistic supernova shock breakout, but the mechanisms are obscure. The breakout of a relativistic shock from a spherical shell of radius R_s would produce a pulse of radiation whose envelope would be proportional to the local (relativistically collimated) emission pattern $f(\theta(t))$, where $\theta(t) = (2ct/R_s)^{1/2}$ and $\theta \sim \gamma^{-1} \ll 1$ is assumed. The rapid fluctuations of intensity under a flat envelope observed in many bursts might then be explained by shock refraction in a turbulent stellar envelope. Unfortunately, the implied values of $R_s \sim 2ct_b\gamma^2$, where t_b is the burst duration, are unreasonably large, particularly because the γ - γ pair-production constraint on 100 MeV gamma-rays implies $\theta < 0.01$.

Such catastrophic events might be expected all to have similar time histories, or to fall into a small number of homogeneous classes, inconsistent with observation. The observed soft gamma repeaters, the 8 second periodicity of the March 5, 1979 burst, the absence of bursts from local supernovae, the identification of 400–500 KeV spectral features with positron annihilation lines of modest redshift, and the complex and varied time histories of intensity within bursts all disfavor these cosmological hypotheses.

If an observer is displaced by a distance a from the center of a sphere of radius $R \gg a$, the mean cosine of the angle θ between the points in the sphere and the direction of displacement is found by integration, keeping only the lowest order terms in a/R :

$$\langle \cos \theta \rangle = \frac{3}{4\pi R^3} \int \int \int d\phi \sin \theta_0 d\theta_0 r^2 dr \cos \theta \approx \frac{a}{R}, \quad (2)$$

where $\theta = \theta_0 + \Delta\theta \approx \theta_0 - \frac{a}{r} \sin \theta_0$ and the integration runs over coordinates (r, θ_0, ϕ)

measured from the center of the sphere. Adopting $a = 8.5$ Kpc for our displacement from the center of the Galaxy and $\langle \cos \theta \rangle < 0.08$ (BATSE 1991; 1σ bound) implies $R > 106$ Kpc; the 3σ bound of $\langle \cos \theta \rangle < 0.22$ relaxes this to $R > 39$ Kpc. There is little other direct evidence for a spherical component of our Galaxy extending to these radii, but it is believed (based on dynamical measurements) that the mass distributions of many galaxies similarly extend far beyond their visible radii. The requirement that the halo be spherical constrains models of its formation and gravitational potential.

It is possible that the objects (by consensus, probably neutron stars) producing gamma-ray bursts were formed in this extended halo early in the history of the Galaxy, and that they remain active for $\geq 10^{10}$ years. The total mean gamma-ray burst flux crossing the Galactic plane is $\sim 10^{-11}$ erg/cm²sec, corresponding to a Galactic luminosity of $\sim 10^{37}$ erg/sec even for a very extended halo ($R \sim 100$ Kpc), or to $\sim 10^{26}$ erg/sec $M_{\odot}$ for a halo mass of $\sim 10^{11} M_{\odot}$, comparable to the mass of the visible Galaxy. The cumulative power radiated is $\sim 3 \times 10^{43}$ erg/ $M_{\odot}$. If all the mass were in neutron stars, this energy would be comparable to their magnetospheric energy for $B \sim 10^{13}$ gauss, or to their rotational energy with moments of inertia of 10^{45} gm cm² and spin periods ~ 25 seconds. Presumably only a fraction of the mass is in neutron stars; their expected magnetospheric energy is inadequate, but their rotational energy may be sufficient.

Alternatively, the sources of gamma-ray bursts may be born within the visible Galaxy but escape to an extended halo. It is well known (Lyne, Anderson and Salter 1982) that many (but probably not all) pulsars are produced with recoil velocities ~ 200 km/sec, sufficient to escape the Galactic disc. When combined with their initial orbital kinetic energy, they may be sufficiently energetic to escape into an extended halo. The time required to fill a halo of radius $R \sim 40$ – 100 Kpc is 1 – 3×10^8 years. This hypothesis is complicated by the fact that their initial orbital velocities about the Galactic center will tend to concentrate the neutron stars to low Galactic latitudes, and any low-recoil fraction will remain in the Galactic plane, concentrated toward the Galactic center.

The lifetimes of these neutron stars as gamma-ray burst sources must not be less than this filling time (or the bursts would be concentrated within the visible Galaxy, and peak towards its center), but could be arbitrarily greater. It is not possible to estimate the number of neutron stars required because their burst frequency is unknown. If the neutron stars freely escape to infinity and radiate indefinitely their $n \propto r^{-2}$ implies an asymptotic $d \log N / d \log S = -0.5$, not as flat as the $d \log N / d \log S = 0$ found for a confined space distribution, but flat enough to explain the data. The energy required is the same as in the case of gamma-ray burst sources born in the halo.

For a simple model of gamma-ray burst sources recoiling from their birthplaces in the Galactic disc, it is possible to calculate the expected dipole moment of their distribution of arrival directions with respect to the Galactic center and the quadrupole moment with respect to the Galactic plane. Assume that we reside a distance r_0 from the center of a Galactic disc of radius L , uniformly populated with source birthplaces, and that each point in the disc uniformly fills with observable gamma-ray bursts a sphere of radius $R \gg L, r_0$ with itself at its center. The uniformly filled sphere is a rough approximation to the motion of neutron stars in the unknown potential of the Galactic halo. Using the result (2), which now describes the observed asymmetry of each sphere, with $a^2 = r^2 + r_0^2 - 2rr_0 \cos \phi$ and (r, ϕ) denoting the coordinates of the disc point, projecting this asymmetry onto the direction to the Galactic center, and integrating over the disc yields

$$\langle \cos \theta \rangle \approx \frac{r_0}{R}, \quad (3)$$

where θ is the angle between a burst's arrival direction and that to the Galactic center. Application to the BATSE data yields the same bounds on R as discussed above, although now R refers to the radius of a sphere populated by gamma-ray bursts from a given neutron star birthplace, rather than the radius of a sphere of source positions centered at the Galactic center; the distinction is slight for $R \gg L, r_0$, as must be the case. If the assumption of uniform filling is replaced by d^{-2} density distribution, where d is the

distance from the birthplace (reflecting an outward streaming distribution, rather than a trapped one), then the result (3) is multiplied by a factor $O(\ln(R/r_0)) \sim 2$.

It is similarly possible to calculate an average $\langle \sin^2 \beta \rangle$, where β is a burst's Galactic latitude. For an observer in the Galactic equatorial plane but at a distance a from the center of a uniformly populated sphere of radius R ,

$$\langle \sin^2 \beta \rangle \approx \frac{1}{3} - \frac{4a^2}{5R^2}, \quad (4)$$

to lowest non-vanishing order in a/R . Averaging over spheres centered in the Galactic disc yields

$$\langle \sin^2 \beta \rangle \approx \frac{1}{3} - \frac{4r_0^2 + 2L^2}{5R^2}. \quad (5)$$

Because of the quadratic dependence on R the resulting bounds are less strict than those obtained from (2) or (3).

The results (3) and (5) may be combined to predict a relation between these two angular averages, if the relation between L and r_0 is known. Take, for simplicity, the reasonable estimate $L = \sqrt{2}r_0$. Then

$$\langle \sin^2 \beta \rangle \approx \frac{1}{3} - \frac{8}{5} \langle \cos \theta \rangle^2. \quad (6)$$

This relation offers a test of the simple geometrical model used here. This test is readily satisfied by the extant BATSE data, but may become more discriminating as more data accumulate.

The gamma-ray burst of March 5, 1979 may, in part, be explained by the extended halo hypothesis. If most other bursts are at distances less than the thickness of the Galactic disc, as was often supposed before the BATSE data became available, then the March 5, 1979 event was extraordinary, because its luminosity must have been $\sim 10^5 - 10^6$ times greater than that of a typical burst. It is hard to reconcile this interpretation with the absence of bursts similar to that of March 5, 1979, but occurring in our own Galaxy, which contains ~ 10 times the mass of the LMC, and might therefore be expected to have ~ 10

times as many such events (but closer and even brighter!). If, instead, bursts in general are distributed throughout a very extended halo of our Galaxy, extending to distances at least as great as that of the LMC, then the March 5, 1979 burst would be explicable as an unusually (but not extraordinarily) luminous member of a class of events with a broad distribution of luminosities.

Another consequence of the hypothesis of the birth of gamma-ray burst sources in the Galactic disc is a further development of the empirical division of neutron star births into high and low recoil classes (Katz 1975). Millisecond pulsars, found in superabundance in dynamically fragile globular clusters (like their probable ancestors, the X-ray binaries) must have been formed recoillessly. Slowly spinning pulsars, most of which are single and are observed (Lyne, Anderson and Salter 1982) to have high space velocities (and hence high recoil) are the ancestors of gamma-ray burst sources.

These categories are nearly disjoint because low-recoil millisecond pulsars retain their companions, initially suppressing their pulsar emission (Shvartsman 1971), but later spinning them up. The observed small magnetic dipole moment μ of millisecond pulsars reduces the power $< \mu^2 c / (3R_{ns}^2)$ potentially released by magnetospheric dissipation in a gamma-ray burst, explaining the absence of this Galactocentrically-concentrated population from the spatial distribution of bursts, although their large rotational energy suggests that a significant luminosity in microbursts may be possible.

High-recoil neutron stars, born with slow spin but which lose any binary companion, must retain their magnetic fields for a very long time in order to produce gamma-ray bursts (empirically, Her X-1 has retained its large field for at least 3×10^7 years, the time required for it to reach its height above the Galactic plane, and perhaps much longer). Low-recoil single neutron stars and high-recoil neutron stars which retain their companions, although individually interesting (Her X-1, PSR1913+16), are uncommon.

If gamma-ray bursts are distant then they must be very powerful. At 100 Kpc a strong 10^{-5} erg/cm² burst, radiating isotropically, emits 10^{43} erg, posing energetic problems

similar to those posed by the March 5, 1979 event (Katz 1982). Nonthermal mechanisms involving the flare-like dissipation of much of a magnetosphere's energy are an attractive explanation of much gamma-ray burst phenomenology, but they are barely energetically adequate for a single burst. The possibility of regeneration of magnetospheric energy even during a burst (perhaps by twisting up the magnetosphere by differential rotation) must be considered. Note, though, that even if the burst population fills a sphere of radius R the brighter bursts will, on average, be significantly closer than R , reducing their energy requirements but increasing the spatial anisotropy of their distribution.

A final consequence of the distances suggested here is that quiescent burst counterparts will be too faint to observe. Stellar companions (unless close and compact) will have been lost when recoil was imparted. Accurate burst coordinates and deep searches will show only blank fields, as has been the case for the few accurate positions presently available (Schaefer 1990). An essentially distance-independent argument (Katz 1985) implies that Schaefer's (1990; but see Żytkow 1990) reported optical flashes can only be explained as nonthermal emission. At a distance of 100 Kpc thermal reprocessing would be inadequate to explain the reported intensities (the inferred absolute magnitudes would be brighter than -15!). Similarly, at these distances self-absorption suppresses the reprocessed X-ray flux, explaining the low X-ray luminosity of gamma-ray bursts (Imamura and Epstein 1987).

I thank G. J. Fishman for supplying data in advance of publication and F. J. Dyson for discussions.

References

- BATSE Science Team 1991, I. A. U. Circ. 5358.
- Cline, T. L. 1984 in *High Energy Transients in Astrophysics*, ed. S. E. Woosley (AIP: New York) p. 333.
- Imamura, J. N., and Epstein, R. I. 1987, *Ap. J.* **313**, 711.
- Jennings, M. C. 1984, in *High Energy Transients in Astrophysics*, ed. S. E. Woosley (AIP: New York) p. 412.
- Katz, J. I. 1975, *Nature* **253**, 698.
- Katz, J. I. 1982, *Ap. J.* **260**, 371.
- Katz, J. I. 1985, *Ap. Lett.* **24**, 183.
- Katz, J. I. 1987, *High Energy Astrophysics* (Addison-Wesley: Menlo Park) §4.5.1.
- Lyne, A. G., Anderson, B., and Salter, M. J. 1982, *M. N. R. A. S.* **201**, 503.
- Meegan, C. A., Fishman, G. J., and Wilson, R. B. 1985, *Ap. J.* **291**, 479.
- Schaefer, B. E. 1990, *Ap. J.* **364**, 590.
- Shvartsman, V. F. 1971, *Sov. Astron.—AJ* **15**, 342.
- Woosley, S. E. 1984, ed. *High Energy Transients in Astrophysics* (AIP: New York).
- Żytkow, A. 1990, *Ap. J.* **359**, 138.